



RESEARCH

Open Access

$\alpha 7$ nAChR/NMDAR coupling affects NMDAR function and object recognition

Shupeng Li^{1†}, Qiang Nai^{1†}, Tatiana V Lipina², John C Roder² and Fang Liu^{1,3*}

Abstract

The $\alpha 7$ nicotinic acetylcholine receptor (nAChR) and NMDA glutamate receptor (NMDAR) are both ligand-gated ion channels permeable to Ca^{2+} and Na^{+} . Previous studies have demonstrated functional modulation of NMDARs by nAChRs, although the molecular mechanism remains largely unknown. We have previously reported that $\alpha 7$ nAChR forms a protein complex with the NMDAR through a protein-protein interaction. We also developed an interfering peptide that is able to disrupt the $\alpha 7$ nAChR-NMDAR complex and blocks cue-induced reinstatement of nicotine-seeking in rat models of relapse. In the present study, we investigated whether the $\alpha 7$ nAChR-NMDAR interaction is responsible for the functional modulation of NMDAR by $\alpha 7$ nAChR using both electrophysiological and behavioral tests. We have found that activation of $\alpha 7$ nAChR upregulates NMDAR-mediated whole cell currents and LTP of mEPSC in cultured hippocampal neurons, which can be abolished by the interfering peptide that disrupts the $\alpha 7$ nAChR-NMDAR interaction. Moreover, administration of the interfering peptide in mice impairs novel object recognition but not Morris water maze performance. Our results suggest that $\alpha 7$ nAChR/NMDAR coupling may selectively affect some aspects of learning and memory.

Keywords: $\alpha 7$ nAChR, NMDAR, Learning, Memory, LTP

Background

Glutamate is the principal excitatory neurotransmitter in brain and N-methyl-D-aspartate (NMDA) receptors, one of the major glutamate receptors, are important in the activity-dependent synaptic plasticity and excitotoxicity that underlies learning, memory, neural development and some neurological disorders [1-3]. Both NMDAR and the $\alpha 7$ nicotinic acetylcholine receptor (nAChR) are ligand-gated ion channel receptors with high Ca^{2+} permeability. NMDARs contain intrinsic ion channels comprised of NR1 subunits, an essential subunit of NMDAR that exists as a number of splice variants, and NR2 subunits, which are encoded by four different gene products, termed NR2A-D [4,5].

Nicotine interacts with nicotinic receptors (nAChRs) in the brain to initiate neuroadaptive changes at both cellular and circuit levels. The nAChRs are composed of five

distinct membrane-spanning subunits (α and β subunits) that combine to form a functional receptor. There are nine isoforms of the neuronal α subunit ($\alpha 2$ - $\beta 10$), and three isoforms of the neuronal β subunit ($\beta 2$ - $\beta 4$) [6]. Various sub-types of nAChRs differ in their subunit composition and sensitivity to nicotine and are expressed in addiction-relevant brain regions including prefrontal cortex, nucleus accumbens, dorsal striatum, and hippocampus [7]. Unlike NMDARs, nAChRs can exist as both heterometric and homo-metric assemblies of these subunits. $\alpha 7$ nAChRs are highly expressed in hippocampus [7].

The activation of nAChRs can modulate glutamatergic neurotransmission in several ways. Previous studies have reported that nicotine facilitates the induction of LTP in the hippocampal CA1 region [8] by the activation of $\alpha 7$ nAChRs on pyramidal cells [9,10]. This induction of LTP can be blocked by AP5, an NMDAR antagonist [11]. Furthermore, *in vivo* nicotine exposure was reported to induce the enhancement of NMDAR currents in the hippocampus [12]. This nicotine effect is maintained during continued nicotine exposure and is accompanied by increased tyrosine phosphorylation of NR2B [13]. In contrast to the presynaptic nAChRs, somatic or postsynaptic

* Correspondence: f.liu.a@utoronto.ca

[†]Equal contributors

¹Department of Neuroscience, Centre for Addiction and Mental Health, Clarke Division, 250 College Street, Toronto, Ontario M5T 1R8, Canada

³Departments of Psychiatry, University of Toronto, Toronto, Ontario M5T 1R8, Canada

Full list of author information is available at the end of the article

nAChRs can initiate a Ca^{2+} signal that can act via calmodulin to reduce the responsiveness of NMDARs, as manifested by evoked excitatory postsynaptic currents (eEPSCs) [14]. Furthermore, NMDAR antagonists have been found to interfere with tolerance, sensitization, physical dependence and conditioning to self-administrated nicotine, as well as other drugs of abuse [15].

We have previously shown that the $\alpha 7$ nAChR interacts with NMDARs and their coupling mediates cue-induced reinstatement of nicotine in rat [16]. In the present study, we plan to investigate the role of $\alpha 7$ nAChR-NMDAR coupling in modulating NMDAR functions. Since both $\alpha 7$ nAChR and NMDAR have been implicated in learning and memory, we will also investigate the behavioral effects of $\alpha 7$ nAChR-NMDAR coupling in some cognitive tests.


Results and discussion

Activation of $\alpha 7$ nAChR increases NMDAR mediated whole-cell currents

Previously, we showed that activation of $\alpha 7$ nAChR by choline facilitates $\alpha 7$ nAChR-NR2A complex formation [16]. To assess the functional impact of the $\alpha 7$ nAChR-NMDAR interaction following $\alpha 7$ nAChR activation, we

examined the effects of $\alpha 7$ nAChR activation on NMDAR-mediated whole-cell currents in rat hippocampal primary cultures. As shown in Figure 1A, co-application of 1 mM choline with 50 μ M NMDA/10 μ M glycine produced a significantly larger current than the current induced by NMDA/Glycine alone (choline/NMDA/glycine: 2036.3 ± 317.2 pA; NMDA/glycine: 812.9 ± 215.5 pA, $n = 43$, $p < 0.05$). The synergistic effect of choline/NMDA co-application is specific to NMDAR since co-application of choline with 100 μ M KA did not enhance whole-cell currents compared to KA treatment alone (Figure 1B).

It is difficult to differentiate whether the observed enhancement of whole-cell current induced by co-application of choline with NMDA is mediated by nicotinic receptors or NMDARs since both receptors are cation ion channel that are permeable to calcium and sodium. However, the observed enhancement of whole cell current induced by co-application of choline with NMDA can be blocked by simultaneous application of the NMDAR channel blocker MK-801 (10 μ M), but not with the nicotinic receptor open channel blocker chlorisondamine (20 μ M) (Figure 1C, D). This suggests that the observed enhancement of whole cell currents is due to ion influx through NMDAR, but not



nicotinic receptors. Furthermore, $\alpha 7$ -nAChR specific antagonists α -bungarotoxin abolish the synergistic effect of choline/NMDA co-application (Figure 1D), indicating that the activation of $\alpha 7$ -nAChR is required for this process.

Activation of $\alpha 7$ nAChR facilitates NMDAR dependent LTP of mEPSCs

To determine whether the $\alpha 7$ nAChR is able to regulate synaptic strength, we examined the miniature excitatory postsynaptic currents (mEPSCs) during LTP upon activation of $\alpha 7$ nAChR. Previous studies have demonstrated that activation of nicotinic acetylcholine receptors facilitates induction of long-term potentiation, although the molecular mechanism underlying this process remains

unknown. Thus, we initiated our investigation by confirming the effect of nicotine on mEPSC during LTP, using the glycine-induced LTP model in rat hippocampal primary neuron cultures. The glycine-induced LTP model is similar to the electrically evoked EPSCs in CA1 neurons in hippocampal slices [17-19]. Consistent with previous studies in brain slices, choline application (1 mM, 8 minutes) significantly enhanced the frequency of mEPSC during LTP produced by glycine application (200 μ M; 3 min) (Figure 2A, C). There is only a small but significant increase in current amplitude mEPSC of LTP (Figure 2B-E), which may reflect the nature of LTP in primary cultures and the recording paradigm [20,21]. We also concluded that the choline-induced upregulation of mEPSC of LTP is NMDAR dependent since D-APV (100 μ M) co-applied




Figure 2 Choline induced upregulation of NMDAR-dependent LTP of mEPSCs in cultured hippocampal neurons. (A) Examples of continuous recordings from individual neurons 5 minutes before (Basal) and 30 minutes after 8-minute stimulation of neurons with 1 mM choline. (B) Single events taken from the basal and choline traces, respectively showing that the amplitude of mEPSCs was increased by choline application. (C) Cumulative fraction plots for mEPSCs inter-event intervals and amplitudes obtained 5 minutes before (Basal) and 30 minutes after choline (8 min, 1 mM). (D) mEPSC amplitudes are normalized to the values from the initial 10 min and plotted over time. Treatment of neurons with choline (8 min, 1 mM) significantly increased the amplitude of the mEPSCs over the time course of recordings; an effect can be abolished by NMDAR antagonist, AP5 (100 μ M). (E) Amplitude histogram summarizes data from groups of individual neurons treated with glycine (200 μ M; 3 min) in the absence or presence of choline (1 mM) or choline/AP5 (100 μ M). Responses obtained 30 min after glycine treatment (26.5 \pm 2.3 pA), 30 min after choline treatment (31.4 \pm 2.7 pA, n=6, *p < 0.01) and 30 minutes after coapplication of choline/APV (25.9 \pm 2.0 pA n=3, **p < 0.05, paired t-test).

with choline blocked the effect of choline on both the frequency (data not shown) and the amplitude (Figure 2E) mEPSC of LTP.

$\alpha 7$ nAChR-NMDA coupling is responsible for modulation of NMDAR function by the activation of $\alpha 7$ nAChR

Next, we determined whether the direct coupling of $\alpha 7$ nAChR-NMDA plays a role in the functional interaction between $\alpha 7$ nAChR and NMDAR. Our previous reports showed $\alpha 7$ nAChR/NMDAR coupling was mediated by a 10 amino acid fragment (L336-M345) within the second intracellular loop of $\alpha 7$ nAChR. Administration of this peptide could disrupt $\alpha 7$ nAChR/NMDAR coupling as shown in the co-immunoprecipitation experiment. Furthermore, this peptide blocked cue-induced nicotine reinstatement in an animal model of relapse [16]. As shown in Figure 3A, B, intracellular application of $\alpha 7$ pep2[L₃₃₆-M₃₄₅] peptide (10 μ M), which has been shown to be able to disrupt $\alpha 7$ nAChR-NMDA coupling, blocked the choline-induced enhancement of NMDA-mediated whole cell currents, while the control peptide, $\alpha 7$ pep1[R₃₁₆-G₃₂₅], has no such effect. These data suggest that the $\alpha 7$ nAChR-NR2A interaction is required for the functional modulation of NMDAR by the activation of $\alpha 7$ nAChR.

Furthermore, we tested the effect of the interfering peptide $\alpha 7$ pep2[L₃₃₆-M₃₄₅] in choline-mediated NMDAR-dependent mEPSC changes during LTP. As shown in Figure 4A-D, intracellular application of $\alpha 7$ pep2[L₃₃₆-M₃₄₅] peptide blocked choline-induced upregulation of mEPSC frequency and amplitude during LTP, indicating

that the $\alpha 7$ nAChR-NR2A interaction is essential for choline-induced modulation of NMDAR-dependent mEPSCs during LTP.

Disruption of the $\alpha 7$ nAChR-NR2A interaction selectively impaired Novel Object Recognition

Both $\alpha 7$ nAChR and NMDARs have been implicated in learning and memory processes. Thus, we sought to investigate whether the $\alpha 7$ nAChR-NR2A interaction might affect learning and memory. We first tested the $\alpha 7$ pep2 [L₃₃₆-M₃₄₅] peptide for possible effects on the Morris water maze. Mice were injected intraperitoneally with TAT- $\alpha 7$ pep2[L₃₃₆-M₃₄₅] (3 ng/g) or TAT- $\alpha 7$ pep1[R₃₁₆-G₃₂₅] 30 min prior to training and probe trials. As shown in Figure 5, there is no difference between $\alpha 7$ pep2[L₃₃₆-M₃₄₅] peptide treated mice and TAT- $\alpha 7$ pep1[R₃₁₆-G₃₂₅] treated mice in latency to find the platform. There is also no difference between the two groups in the acquisition phase, nor in the probe trial, indicating that the disruption of the $\alpha 7$ nAChR-NMDAR interaction has no effect on the spatial learning and memory required for this task.

To further evaluate the effect of our interfering peptide on cognition, we used two other behavioral tests: the displaced object recognition task and the novel object recognition task. As shown in Figure 6 A, TAT- $\alpha 7$ pep2 [L₃₃₆-M₃₄₅] peptide treatment, but not TAT- $\alpha 7$ pep1[R₃₁₆-G₃₂₅] treatment, induces impairment in novel object recognition in mice. In contrast, there is no difference between the two groups in the displaced object recognition task (Figure 6B). To investigate whether the TAT- $\alpha 7$ pep2[L₃₃₆-M₃₄₅] peptide might affect anxiety-




Figure 3 Application of $\alpha 7$ pep2 peptide blocked choline induced upregulation of NMDA current in hippocampal primary culture. (A) The choline-induced synergistic effect is significantly inhibited by intracellular application of interfering peptides $\alpha 7$ pep2, but not $\alpha 7$ pep1 (10 μ M) (choline/NMDA: 1202.7 ± 182.1 pA; NMDA: 910.5 ± 130.8 , $n = 6$, $p > 0.05$). Cells were held at -70 mV, 20 mM bicuculline, 1 mM strychnine, 0.5 μ M TTX, 1 mM glycine were included in the extracellular solution. **(B)** Amplitude histogram summarizes data from groups of individual neurons treated with glycine (200 μ M; 3 min) in the absence or presence of choline (1 mM) with the intracellular application of $\alpha 7$ pep1, $\alpha 7$ pep2 peptide respectively. Responses obtained 30 min after glycine treatment (basal) and 30 min after choline treatment (choline). $\alpha 7$ pep1 peptide did not block the enhancing effect of choline on the mEPSC amplitude (basal: 25.2 ± 2.1 pA; choline: 28.4 ± 2.4 pA, $n = 4$, $*p < 0.01$, paired t-test) while choline failed to upregulate mEPSC amplitude with the presence of $\alpha 7$ pep2 (basal: 24.2 ± 2.0 ; choline: 25.1 ± 2.3 pA, $n = 6$, $p > 0.05$, paired t-test).




Figure 4 Application of $\alpha 7$ pep2 peptide blocked choline induced upregulation of mEPSC of LTP in hippocampal primary culture.

(A) Examples of continuous recordings from individual neurons 40 minute after intracellular application of $\alpha 7$ pep2 peptide (10 μ M) with/without the presence of choline (1 mM, 8 min). (B) Single events taken from the basal and choline traces after intracellular application of $\alpha 7$ pep2 peptide, showing that choline application failed to increase the amplitude of mEPSCs. (C) Cumulative fraction plots for mEPSCs inter-event intervals and amplitudes obtained 5 minutes before (Basal) and 30 minutes after choline (8 min, 1 mM) with the presence of $\alpha 7$ pep2 peptide intracellularly. (D) Amplitude histogram summarizes data from groups of individual neurons treated with glycine (200 μ M; 3 min) in the absence or presence of choline (1 mM) with the intracellular application of $\alpha 7$ pep1, $\alpha 7$ pep2 peptide respectively. Responses obtained 30 min after glycine treatment (basal) and 30 min after choline treatment (choline). $\alpha 7$ pep1 peptide did not block the enhancing effect of choline on the mEPSC amplitude (basal: 25.2 \pm 2.1 pA; choline: 28.4 \pm 2.4 pA, n = 4, *p < 0.01, paired t-test) while choline failed to upregulate mEPSC amplitude with the presence of $\alpha 7$ pep2 (basal: 24.2 \pm 2.0; choline: 25.1 \pm 2.3 pA, n = 6, p > 0.05, paired t-test).

related behaviour, we tested the effect of TAT- $\alpha 7$ pep2 [L₃₃₆-M₃₄₅] in the elevated plus maze. As shown in Figure 6C, there is no difference between the two groups in the number of entries into the open arms, the time spent on the open arms and the head dips. There is also no difference between the two groups in the total distance travelled, margin and central distance travelled, and time spent in the marginal and central zones (Additional file 1: Figure S1A-E). Taken together, our findings suggest that the $\alpha 7$ nAChR-NMDAR interaction may selectively impair novel object recognition.

In the present study, we provide evidence that the $\alpha 7$ nACh-NMDAR complex modulates NMDAR-mediated whole cell currents and LTP. Furthermore, disruption of this complex via an interfering protein peptide TAT- $\alpha 7$ pep2 [L₃₃₆-M₃₄₅] had no effect on Morris water maze and displaced object recognition in mice, but specifically impaired novel object recognition. Our study

provide the first demonstration that $\alpha 7$ nAChR regulates NMDA-mediated whole cell currents and LTP through a protein-protein interaction. More interestingly, our data suggest that the $\alpha 7$ nAChR-NR2A interaction may specifically play a role in non-spatial learning and memory.

Regulation of ligand-gated ion channel function was traditionally thought to be mainly regulated by receptor phosphorylation and trafficking [22,23]. Receptor phosphorylation involves intracellular second-messengers, including various phosphatase/kinases, while receptor trafficking can be induced by either receptor phosphorylation or direct coupling with intracellular proteins that lead to changes in receptor conformation or receptor plasma membrane expression. Thus, receptor phosphorylation, conformational changes and plasma membrane expression constitute the major means to modulate ligand-gated ion channel function [24].




Figure 5 TAT- α 7pep2 peptide treatment has no effects on spatial learning and memory. Latency to find the platform of mice was not affected by peptide treatment in the Morris water maze task. **(A)** In the acquisition phase, escape latency to find a hidden platform located in the southeast (SE) quadrant was unaffected by treatment. **(B)** Histogram of percent time spent in each quadrant at probe test.

We did not directly investigate the mechanism by which the α 7nAChR-NR2A interaction can lead to enhanced NMDA currents. However, we speculate that there are several possible mechanisms: enhanced phosphorylation, conformational changes or altered cell surface expression of NMDAR. Currently, there is no evidence supporting that either α 7nAChR or NMDAR are able to directly activate second messenger systems, however, both receptors are calcium permeable [25]. It is possible that the calcium influx induced by the activation of α 7nAChR may lead to the activation of intracellular signaling pathways that lead to changes in NMDAR phosphorylation, with potential downstream functional changes. The α 7nAChR-NMDAR complex may also induce conformational changes in the NMDAR or enhance NMDAR cell surface expression that could alter current flow.

We have found that the administration of the interfering peptide in mice impairs novel object recognition, but not Morris water maze performance and displaced object recognition, suggesting that the α 7nAChR-NR2A interaction may specifically play a role in non-spatial learning and memory. However, the fact that our interfering

peptide blocks choline-induced upregulation of mEPSC during LTP, which contributes to spatial learning performance, seems contradictory. LTP has been recognized as a cellular model for learning and memory. Although LTP is generally thought to be associated with spatial learning and memory, there are examples of inconsistencies between LTP and Morris water maze performance, a common way of testing spatial learning and memory. For example, Dr. Morris' lab reported in 1995 [26] that AP5 (an NMDA receptor antagonist) impaired both LTP and water maze performance. However, the AP5-induced learning deficit in the water maze can be prevented if rats are pre-trained in a different water maze before administration of AP5, implying that NMDA receptor-mediated LTP may not be required for all components of spatial learning. In addition, the same issue of Nature includes another paper by Saucier and Cain that shows NMDA receptor-mediated dentate LTP is not required for normal spatial learning in the water maze [27]. Thus, our result showing that disruption of α 7nAChR-NR2A interaction impairs LTP, but not Morris water maze performance, seems contradictory, but it is not unique.




Figure 6 TAT- α 7pep2 Peptide treatment affected nonspatial learning and memory. TAT- α 7pep2 Peptide treatment impaires novel object recognition (A) but not displaced object recognition (B). Times of exploration of the DO and NDO were recorded and expressed as a percentage of the total time of objects investigated. In the novel object recognition session, one of the familiar NDOs were replaced with a new object (NO) at the same location and the two familiar DOs were removed. Data were analyzed with ANOVA with treatment as a between-subjects factor, and object rearrangement or object replacement as a repeated measures factor. The Tukey test was used for post hoc comparisons when ANOVA yielded statistically significant main effects or interactions. (C) In the elevated plus maze, no significant changes in the percent of time spent in open arms, entries into the open arms and head dips were observed in different treatment groups.

We have previously found that disruption of $\alpha 7$ nAChR-NMDAR complex using TAT- $\alpha 7$ pep2[L₃₃₆-M₃₄₅] blocks cue induced reinstatement of nicotine self-administration in rats [16]. In the current study, we have further explored behavioral changes in mice given TAT- $\alpha 7$ pep2[L₃₃₆-M₃₄₅]. We found that TAT- $\alpha 7$ pep2[L₃₃₆-M₃₄₅] had no effect on spatial learning and memory in the Morris water maze and displaced object recognition task in mice, but did affect novel object recognition. This is consistent with previous findings that systemic administration of selective $\alpha 7$ nAChR agonists reverse working memory impairments caused by NMDAR blockade in several behavioral tasks including the 16-arm radial maze, Y-maze, Morris water maze and linear maze, and novel object recognition test [28-31]. There are also other examples of functional interaction between the $\alpha 7$ nAChR and NMDAR. Cholinergic innervation of the hippocampus modulates activity-dependent synaptic plasticity, such as long-term potentiation (LTP) and other processes that contribute to learning and memory [32]. Nicotine was found to enhance LTP of EPSPs in the dentate gyrus and to convert weak stimuli-evoked short-term potentiation into LTP in the CA1. The selective $\alpha 7$ nAChR agonists choline and 2,4-dimethoxybenzylidene anabaseine have also been found to mimic the facilitative action of nicotine in potentiating LTP [33-35], although the mechanisms underlying the effects of $\alpha 7$ nAChR on NMDAR-mediated function remain unclear.

Conclusions

Our results confirm a physical interaction between $\alpha 7$ nAChR and the NR2A of NMDAR that affects both NMDAR-mediated function and novel object recognition. These findings increase our understanding of these two receptor systems and suggest future experiments to further investigate the mechanisms underlying the functional effects of the interaction.

Methods

Primary cultures of dissociated cells

Hippocampi were collected from fetal (E18) Wistar rats. Fetuses are removed from pregnant rats anesthetized by inhalation of isoflurane and killed by cervical dislocation. The dissection and dissociation were performed in ice-cold Hank's balanced salt solution (HBSS, without Ca⁺⁺ and Mg⁺⁺ Gibco) supplemented with 10 mM HEPES (pH 7.4) and 1 mM sodium pyruvate. Neurons were mechanically dispersed by trituration using glass Pasteur pipettes with reduced tips and then added to plating solution composed of 89.5% Neural Basal (NB), 10% horse serum, and 0.5% Penicillin/streptomycin (P/S) [36]. The cells were plated on German origin glass coverslips coated with 0.1 mg/ml poly-d-lysine in Borate Buffer. The cell density was about 50,000-80,000/ml. After 5/6 hours of plating, half of the plating solution was replaced by feeding

solution containing 98% NB, 2% B-27 supplement, 0.5 mM L-glutamine and 0.5% P/S (all from Gibco). The cultures were maintained by feeding twice weekly by replacing half of the solution with fresh feeding solution. After 6 days of plating, 5 μ M Ara-C was added to stop the growth of glial cells.

Electrophysiology

Miniature excitatory postsynaptic currents (mEPSCs) were recorded from cultured hippocampal neurons 2 to 4 weeks days after plating under a whole-cell patch clamp configuration [36]. Electrodes (3-5 M Ω) were pulled from high lead pipettes (Corning 8161, Warner Instruments). Cells were voltage clamped at -70 mV. Access resistance is below 10 M Ω ; recordings with access resistance varying more than 10% were rejected from analysis. The extracellular solution contained (in mM) NaCl 140, CaCl₂ 1.3, KCl 5.0, HEPES 25, glucose 33, TTX 0.0005, strychnine 0.001, and bicuculline methiodide 0.01, at pH 7.4 and osmolarity 325-335 mosmol⁻¹. Each of the tested cells was continuously perfused with the extracellular solution from a single barrel of a computer-controlled multi-barreled fast-step perfusion system (Warner Instruments Inc.). The receptor agonists were applied from different barrel(s).

The response to nicotinic agonists by different hippocampal cultures was variable. Overall, about 30% (53 of 170 cells) of the cells displayed positive nicotinic responses (more than ten times the basal RMS noise level). Only responsive cells were used for further whole-cell or synaptic activity analysis. The intracellular solution consisted of (in mM): CsCl₂ 140, EGTA 2.5, MgCl₂ 2, HEPES 10, TEA 2, and K₂ATP 4, at pH 7.3; and osmolarity 300 to 310 mosmol⁻¹ [19]. In some experiments 10 μ M of peptides $\alpha 7$ pep1 and $\alpha 7$ pep2 were included in the intracellular solution and dialyzed for 30 minutes before recording. Recordings were made at room temperature (21-23°C). Series resistance was not compensated. Synaptic activity was recorded using an Axopatch 200B (Axon Instruments, Inc.); signals were filtered at 2 kHz, digitized at 10 kHz, and stored in a lab computer. Data were analyzed using Mini Analysis Software (Synaptosoft, Inc.). mEPSC frequency and amplitude for each time point were obtained from a two minute recording. The trigger level for event detection was three times higher than that of baseline noise. Visual inspection was performed to eliminate false events. Data were expressed as mean \pm SEM, t-test were used to test the statistical significance of differences between groups.

Behavioural testing

All animal procedures were conducted in accordance with the requirements of the Province of Ontario Animals for Research Act, 1971 and the Canadian Council on Animal

Care (CCAC 1984, 1995). To examine the effects of $\alpha 7$ pep2[L₃₃₆-M₃₄₅] in learning and memory, C57BL/6 J mice were used for the water maze and object recognition tasks. In the water maze task, a single daily intraperitoneal injection of $\alpha 7$ pep2[L₃₃₆-M₃₄₅] (3 ng/g) or vehicle was administered 30 min prior to training and on the probe trial day. For the object recognition task, a single intraperitoneal injection of $\alpha 7$ pep2 or vehicle was administered 30 min prior to object recognition testing.

Locomotor activity was monitored in a directly illuminated (600 lux) clear Perspex chamber (42 × 42 × 30 cm; Accuscan Instruments Inc., Columbus, OH, USA) by interruptions of 16 horizontal and 16 vertical sensors (infrared beams) spaced 2.5 cm apart. Data was recorded every 5 min of the testing period. Data were analyzed with two way analyses of variance (anovas) with treatment as main factor and repeated measures (time intervals).

For the Morris water maze task, 12–16-wk-old C57BL/6 J mice were used. The water maze consisted of a 185 cm diameter cylindrical tank containing a 15 cm circular platform and water (26 ± 1°C) rendered opaque by the addition of white non-toxic paint. The training regime consisted of acquisition training to a hidden platform in the southeast (SE) quadrant for 3 d (day 1–3; six trials per day; maximum duration, 90 s; ITI, 40 min). Probe trials (90 s duration) were administered 18 h after the last acquisition. All behavioural events were video recorded and analyzed using Observer 5.0 software (Noldus Information Technology). Behavioural data for escape latency were analyzed using a two-way ANOVA with training days as repeated measurement. For the probe trials, statistical comparisons between groups for the time over quadrants were done using one-way ANOVA with the critical α level set to 0.05 for all statistical analyses.

Object recognition tests were performed as described [37] using a modified open field set-up. The open field apparatus consisted of a square box (41 × 41 × 33 cm) made of clear Perspex (Ugo Basile) that was connected to horizontal and vertical infrared sensors. During the habituation session, four different plastic objects were presented in the open field: a cube (5 × 5 × 5 cm), hollow cylinder (6 cm height and 4 cm diameter), solid cylinder (3 cm height × 6 cm diameter), and prism (3.5 × 4.5 × 6 cm). Exploration of the four different plastic objects in the open field were measured every 5 min for 15 min under dim lighting (habituation).

In the displace object recognition session, the four objects, initially placed in a square arrangement, were reconfigured into a polygon-shaped pattern by moving two objects (the displaced objects or DO). The remaining two objects were left at the same location (nondisplaced objects [NDOs]). Times of exploration of the DO and NDO were recorded for 5 min and expressed as a percentage of the total time spent investigating objects. In the novel

object recognition session, one of the familiar NDOs was replaced with a new object (NO) at the same location and the two familiar DOs were removed. The time examining a NO or a familiar object (FO) was recorded for 5 min and was expressed as a percentage of the total time spent investigating objects. Data were analyzed using ANOVA with drug treatment as a between-subjects factor, and object rearrangement or object replacement as a repeated measures factor. The Tukey test was used for post hoc comparisons when the ANOVA yielded statistically significant main effects or interactions.

Elevated plus maze (EPM)

Experiments were conducted in a dimly lit room with a light intensity on the central platform of 210 lux [38]. During a 5-min observation period, the number of entries (defined as four paws into a maze arm) and the amount of time spent in the open arms, closed arms and the central platform were scored by the observer. The total number of entries for each subject was collected. These data are presented as percentage time spent in closed or open arm/total duration of experiment × 100. Data were analyzed using ANOVA with drug treatment as a between-subjects factor.

Additional file

Additional file 1: Supplemental Figures TAT- $\alpha 7$ pep2 peptide treatment has no effects on locomotor activity. TAT- $\alpha 7$ pep2 peptide treatment did not affect total distance travelled (Supplemental Figure 1A), Margin Distance Travelled (Supplemental Figure 1B), Margin Time Spent (Supplemental Figure 1C), Centre Distance Travelled (Supplemental Figure 1D), Centre Time Spent (Supplemental Figure 1E) in the open field test.

Abbreviations

EPM: Elevated plus maze; DO: Displaced objects; NDOs: Nondisplaced objects; NO: New object; FO: Familiar object; mEPSCs: Miniature excitatory postsynaptic currents; LTP: Long-term potentiation; eEPSCs: Evoked excitatory postsynaptic currents; nAChR: nicotinic acetylcholine receptor; NMDAR: N-methyl-D-aspartate receptor.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SL and TL conducted the behavior tests and data analysis; QN conducted the electrophysiology experiments and analyzed all the electrophysiology data. JR advised the design of the behavior tests. FL designed, supervised the study and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by operating grant from Canadian Institute of Health Research to FL.

Author details

¹Department of Neuroscience, Centre for Addiction and Mental Health, Clarke Division, 250 College Street, Toronto, Ontario M5T 1R8, Canada.

²Samuel Lunenfeld Research Institute at Mount Sinai Hospital, Toronto, Ontario M5T 1R8, Canada. ³Departments of Psychiatry, University of Toronto, Toronto, Ontario M5T 1R8, Canada.

Received: 3 October 2013 Accepted: 12 December 2013
Published: 20 December 2013

References

- Bartlett TE, Wang YT: The intersections of NMDAR-dependent synaptic plasticity and cell survival. *Neuropharmacology* 2013, **74**:59–68.
- Bliss TV, Collingridge GL: Expression of NMDA receptor-dependent LTP in the hippocampus: bridging the divide. *Mol Brain* 2013, **6**:5.
- Paoletti P, Bellone C, Zhou Q: NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease. *Nat Rev Neurosci* 2013, **14**(6):383–400.
- Gray R, Rajan AS, Radcliffe KA, Yakehiro M, Dani JA: Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature* 1996, **383**(6602):713–716.
- MacDermott AB, Mayer ML, Westbrook GL, Smith SJ, Barker JL: NMDA-receptor activation increases cytoplasmic calcium concentration in cultured spinal cord neurones. *Nature* 1986, **321**(6069):519–522.
- Changeux JP, Edelman SJ: Allosteric mechanisms of signal transduction. *Science* 2005, **308**(5727):1424–1428.
- Sargent PB: The diversity of neuronal nicotinic acetylcholine receptors. *Annu Rev Neurosci* 1993, **16**:403–443.
- Nakauchi S, Sumikawa K: Endogenously released ACh and exogenous nicotine differentially facilitate long-term potentiation induction in the hippocampal CA1 region of mice. *Eur J Neurosci* 2012, **35**(9):1381–1395.
- Soderman A, Mikkelsen JD, West MJ, Christensen DZ, Jensen MS: Activation of nicotinic alpha(7) acetylcholine receptor enhances long term potentiation in wild type mice but not in APP(swe)/PS1DeltaE9 mice. *Neurosci Lett* 2011, **487**(3):325–329.
- Lagostena L, Trocme-Thibierge C, Morain P, Cherubini E: The partial alpha7 nicotine acetylcholine receptor agonist S 24795 enhances long-term potentiation at CA3-CA1 synapses in the adult mouse hippocampus. *Neuropharmacology* 2008, **54**(4):676–685.
- Mann EO, Greenfield SA: Novel modulatory mechanisms revealed by the sustained application of nicotine in the guinea-pig hippocampus in vitro. *J Physiol* 2003, **551**(Pt 2):539–550.
- Yamazaki Y, Jia Y, Niu R, Sumikawa K: Nicotine exposure in vivo induces long-lasting enhancement of NMDA receptor-mediated currents in the hippocampus. *Eur J Neurosci* 2006, **23**(7):1819–1828.
- Nakajima A, Kinugasa Y, Torii J, Hishinuma T, Tomioka Y, Yamada K, Yamakuni T: Repeated treatment with nicotine induces phosphorylation of NMDA receptor NR2B subunit in the brain regions involved in behavioral sensitization. *Neurosci Lett* 2012, **524**(2):133–138.
- Fisher JL, Dani JA: Nicotinic receptors on hippocampal cultures can increase synaptic glutamate currents while decreasing the NMDA-receptor component. *Neuropharmacology* 2000, **39**(13):2756–2769.
- Trujillo KA, Akil H: Excitatory amino acids and drugs of abuse: a role for N-methyl-D-aspartate receptors in drug tolerance, sensitization and physical dependence. *Drug Alcohol Depend* 1995, **38**(2):139–154.
- Li S, Li Z, Pei L, Le AD, Liu F: The alpha7nACh-NMDA receptor complex is involved in cue-induced reinstatement of nicotine seeking. *J Exp Med* 2012, **209**(12):2141–2147.
- Liao D, Hessler NA, Malinow R: Activation of postsynaptically silent synapses during pairing-induced LTP in CA1 region of hippocampal slice. *Nature* 1995, **375**(6530):400–404.
- Lu W, Man H, Ju W, Trimble WS, MacDonald JF, Wang YT: Activation of synaptic NMDA receptors induces membrane insertion of new AMPA receptors and LTP in cultured hippocampal neurons. *Neuron* 2001, **29**(1):243–254.
- Man HY, Wang Q, Lu WY, Ju W, Ahmadian G, Liu L, D'Souza S, Wong TP, Taghibiglou C, Lu J, et al: Activation of PI3-kinase is required for AMPA receptor insertion during LTP of mEPSCs in cultured hippocampal neurons. *Neuron* 2003, **38**(4):611–624.
- Fitzjohn SM, Pickard L, Duckworth JK, Molnar E, Henley JM, Collingridge GL, Noel J: An electrophysiological characterisation of long-term potentiation in cultured dissociated hippocampal neurones. *Neuropharmacology* 2001, **41**(6):693–699.
- Sokolova IV, Lester HA, Davidson N: Postsynaptic mechanisms are essential for forskolin-induced potentiation of synaptic transmission. *J Neurophysiol* 2006, **95**(4):2570–2579.
- Swope SL, Moss SJ, Raymond LA, Huganir RL: Regulation of ligand-gated ion channels by protein phosphorylation. *Adv Second Messenger Phosphoprotein Res* 1999, **33**:49–78.
- Schwappach B: An overview of trafficking and assembly of neurotransmitter receptors and ion channels (Review). *Mol Membr Biol* 2008, **25**(4):270–278.
- Mayer ML: Glutamate receptor ion channels. *Curr Opin Neurobiol* 2005, **15**(3):282–288.
- Uteshev VV: Evaluation of Ca2+ permeability of nicotinic acetylcholine receptors in hypothalamic histaminergic neurons. *Acta Biochim Biophys Sin (Shanghai)* 2010, **42**(1):8–20.
- Bannerman DM, Good MA, Butcher SP, Ramsay M, Morris RG: Distinct components of spatial learning revealed by prior training and NMDA receptor blockade. *Nature* 1995, **378**(6553):182–186.
- Saucier D, Cain DP: Spatial learning without NMDA receptor-dependent long-term potentiation. *Nature* 1995, **378**(6553):186–189.
- Pichat P, Bergis OE, Terranova JP, Urani A, Duarte C, Santucci V, Gueudet C, Voltz C, Steinberg R, Stemmelin J, et al: SSR180711, a novel selective alpha7 nicotinic receptor partial agonist: (II) efficacy in experimental models predictive of activity against cognitive symptoms of schizophrenia. *Neuropsychopharmacology* 2007, **32**(1):17–34.
- Roncarati R, Scali C, Comery TA, Grauer SM, Aschmi S, Bothmann H, Jow B, Kowal D, Gianfriddo M, Kelley C, et al: Procognitive and neuroprotective activity of a novel alpha7 nicotinic acetylcholine receptor agonist for treatment of neurodegenerative and cognitive disorders. *J Pharmacol Exp Ther* 2009, **329**(2):459–468.
- Arendash GW, Sengstock GJ, Sanberg PR, Kem WR: Improved learning and memory in aged rats with chronic administration of the nicotinic receptor agonist GTS-21. *Brain Res* 1995, **674**(2):252–259.
- Marighetto A, Valerio S, Desmedt A, Philippin JN, Trocme-Thibierge C, Morain P: Comparative effects of the alpha7 nicotinic partial agonist, S 24795, and the cholinesterase inhibitor, donepezil, against aging-related deficits in declarative and working memory in mice. *Psychopharmacology (Berl)* 2008, **197**(3):499–508.
- Sugisaki E, Fukushima Y, Tsukada M, Aihara T: Cholinergic modulation on spike timing-dependent plasticity in hippocampal CA1 network. *Neuroscience* 2011, **192**:91–101.
- Matsuyama S, Matsumoto A, Enomoto T, Nishizaki T: Activation of nicotinic acetylcholine receptors induces long-term potentiation in vivo in the intact mouse dentate gyrus. *Eur J Neurosci* 2000, **12**(10):3741–3747.
- Nishimoto T, Kadoyama K, Taniguchi T, Takano M, Otani M, Nakamura-Hirota T, Lu Y, Matsumoto A, Matsuyama S: Synaptotagmin1 synthesis induced by synaptic plasticity in mouse hippocampus through activation of nicotinic acetylcholine receptors. *Neurosci Lett* 2011, **489**(1):25–29.
- Chen L, Yamada K, Nabeshima T, Sokabe M: alpha7 Nicotinic acetylcholine receptor as a target to rescue deficit in hippocampal LTP induction in beta-amyloid infused rats. *Neuropharmacology* 2006, **50**(2):254–268.
- Nai Q, Li S, Wang SH, Liu J, Lee FJ, Frankland PW, Liu F: Uncoupling the D1-N-methyl-D-aspartate (NMDA) receptor complex promotes NMDA-dependent long-term potentiation and working memory. *Biol Psychiatry* 2010, **67**(3):246–254.
- Ng D, Pitcher GM, Szilard RK, Sertie A, Kanisek M, Clapcote SJ, Lipina T, Kalia LV, Joo D, McKerlie C, et al: Neto1 is a novel CUB-domain NMDA receptor-interacting protein required for synaptic plasticity and learning. *PLoS Biol* 2009, **7**(2):e41.
- Haque FN, Lipina TV, Roder JC, Wong AH: Social defeat interacts with Disc1 mutations in the mouse to affect behavior. *Behav Brain Res* 2012, **233**(2):337–344.

doi:10.1186/1756-6606-6-58

Cite this article as: Li et al.: a7nAChR/NMDAR coupling affects NMDAR function and object recognition. *Molecular Brain* 2013 **6**:58.